

# PLANT DISEASE AND INSECT ADVISORY



Department Entomology and Plant Pathology  
Oklahoma State University  
127 Noble Research Center  
Stillwater, OK 74078



Vol. 1, No. 24

Website: <http://plants.okstate.edu/Pddl/advisory.htm>

Aug 14, 2002

## Preparing for Pecan Weevil Season and Suggestions for use of Circle Traps


Phil Mulder, Extension Entomologist



With most of the early season insect problems behind us it is time to pay more attention to pecan weevil season. Now is the time to prepare for the upcoming barrage of weevils that can infest the crop. With the recent rains experienced across most the state, pecan weevils have begun to emerge in several locations, particularly from sandier soils. September is the prime time for pecan weevil activity, with the third week generally representing the peak period. Early damage from pecan weevil can vary from year to

year and orchard to orchard. In fact, susceptibility to this pest can vary from tree to tree in a single, home yard situation. In years past, reports have surfaced of 75% crop loss due to pecan weevil injury. In addition to the injury on the crop itself, buyers and processors can purchase affected pecans and have to either clean them out, or dispose of the crop. In the native pecan growing range, this pest has evolved with the host and is common throughout much of the production areas of Texas, Oklahoma, and Georgia. In recent years; however, the pecan weevil found its way into the "western irrigated area." Through all the efforts of the growers, state agriculture departments, and scientists, much of the New Mexico area still remains weevil free. I'm sure it will be an ongoing and occasionally exhaustive effort to keep the pecan weevil from establishing a foothold in that part of the country, as they have witnessed its encroachment twice in the last three years.

The typical life cycle of the pecan weevil requires 2 to 3 years, with the majority of weevils emerging two years after burrowing into the soil. Once a weevil larva enters the soil, it has likely matured through four instars. The final instar (fourth) may feed for a short while (about one week) but generally spends the one to two years in the soil without feeding. Larvae will burrow into the soil at various depths depending on soil type, moisture and condition of the orchard floor. Most larvae are likely 6-9 inches deep in the soil profile. Ninety percent of the larvae that entered the soil on "year 1" will pupate in "year 2." The remainder of the population will not pupate until "year 3." The pupal stage lasts for about 3-4 weeks (longer in females than in males) and the adults then remain inactive within their earthen cells and do not attempt



emergence from the soil until “year 3” (about 2 years after they entered the soil as larvae). As stated previously, about 10% of the population will not emerge until “year 4.” After emergence from the soil, adult weevils live for about 2 to 3 weeks, with females surviving longer than males. Female weevils are capable of surviving much longer if nuts are not acceptable for oviposition (egg-laying). If the nuts are not ready for oviposition by female weevils, then adults will cause feeding damage on the nuts. This damage may consist of feeding punctures, prior to shell hardening, which will cause premature drop of the pecan. Feeding by pecan weevil after shell hardening may cause the deterioration of the gel resulting in “sticktights.” If the weevil penetrates only the shuck, then damage may be limited to slight scars, black spots, pits or molds on the kernel near the puncture site.

Oviposition by female pecan weevils can occur as early as 2 days after emergence; however, the majority of egg production occurs 10-12 days after emergence. Regardless of how soon oviposition begins, it is initiated on early maturing varieties sooner than on trees that have late maturing nuts. Each female can average around 35-55 eggs deposited during her life. She will average about four eggs per nut. With a healthy weevil population, this can account for a great amount of damage and of course a tremendous carry-over in the weevil population for subsequent years.



Much of the damage experienced by pecan weevil can be avoided with timely use of insecticides. Making the decision on when to treat; however, can be a dilemma for many growers. That decision should be made with some sort of trapping program established in the orchard. For trapping we suggest the pecan weevil Circle trap. Although growers have become familiar with these traps over the last few years, many questions remain on how to best utilize the traps. Several studies conducted in Oklahoma over the last four years have helped

provide some answers to many of these questions. Following are several suggestions on the utility of the Circle trap and methods for arriving at a treatment threshold based on captures.

- 1) Commercial situations (several trees) – Use as many as you can afford, to represent the land area, soil types, pecan varieties, and topography in your area.
- 2) Where possible, target indicator trees – Those with a history of weevil problems, those that are earlier than most, or those located in sandy soils.
- 3) Place two traps on each targeted tree. One on the north, west or northwest side of the tree and the other just opposite. Set them at a height suitable for your operation. If cattle are going to be grazing the orchard floor, then put the traps sufficiently high on the trunk to avoid their natural curiosity. Just remember that someone needs to check them regularly, therefore, they should be reachable for the person doing the checking. Where cattle are present, a small step ladder or truck pickup bed will help. Soon after placement on the trees, cattle lose their curiosity with these traps, probably for two reasons. First they're made of aluminum screening, which they don't prefer to chew on and second because after a few days they tend to recognize the trap as part of the tree.
- 4) When attempting to encircle the tree trunk, place additional wire screening wherever the traps don't match up, and curl the top lip of the screening over to funnel the weevils up into the trap tops.
- 5) Mow grass under the trees to avoid providing weevils a spot to take off into the canopy.
- 6) Begin trapping by July 15 if possible, but don't be overly concerned if the nuts are still in the water stage.
- 7) Do not cease trapping too early (before mid-October).
- 8) Inspect traps at least every 2 to 3 days. Set up a M-W-F checking schedule, and stick to it.
- 9) Keep a running set of written records, to include date of trap inspection, number of weevils caught, sex of weevils caught, and tree number or location. This information should be retained for long term records and can help you identify indicator spots and/or trees.
- 10) Convert the total number of weevils collected from all traps on each inspection date into the number of weevils/trap/day. In other words, the total number of weevils from "X" traps after "X" number of days.



- EXAMPLE: You examined 16 traps on 8 trees on Monday and found nothing but on Wednesday you gathered up 12 weevils from the same 16 traps. Therefore, divide 12 by (16 times 2). The two represents the days since your last check of the traps.  $12/32 = 0.375$  weevils/trap/day.
- 11) Begin control measures if the number of weevils equals or exceeds 0.3 weevils/trap/day.
  - 12) Be sure to regularly clean the trap screen tops to remove spiders, lacewings, webs and dead insects which may interfere with functioning of the trap. A stiff toothbrush can help in this task.
  - 13) Be diligent in this task from July to October, but really focus immediately following any 1"-2" rainfall. If conditions are hot and dry, as they often are in July and August, weevil emergence may be delayed but can still be significant up until shuck-split. In Oklahoma, populations generally peak about the third week of September.
  - 14) When an application is about to be made, remove the top cylinder of each trap to allow everything to crawl through. The air-blast sprayers and the insecticide will cause a great

many weevils to fall shortly after treatment. These "sick weevils" will begin crawling up tree trunks and entering traps. This can give the impression of an ineffective treatment or that you had a fresh emergence. Leave tops off for a period of one week, then resume the monitoring program.

For homeowners that may have one or a few trees, it may be possible to "trap-out" the majority of their weevil population by encircling the trees with traps and following the same suggestions. This may reduce the number of insecticide applications needed in this sensitive environment and yet preserve pecans for harvesting. In localized "hot pockets," where weevils have relied on a few trees in the area, thorough monitoring and treatment over a two to three year period could dramatically reduce or even eliminate the pecan weevil for several years.

Information on construction of the Circle traps and their utility can be obtained from OSU Extension Fact Sheet 7190 "Monitoring adult weevil populations in pecan and fruit trees in Oklahoma." Several vendors can supply the traps for you and they range in cost from \$10 to \$25 per trap. For a list of potential vendors you can simply contact my office.

---

Oklahoma State University, in compliance with Title IV and VII of the Civil Rights Act of 1964, Executive Order of 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, sex, age, religion, disability, or status as a veteran in any of its policies, practices or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Samuel E. Curl, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Dean of Agricultural Sciences and Natural Resources.